

Non-uniform ground state for the Bose gas

András Sütő

Research Institute for Solid State Physics

Hungarian Academy of Sciences

P. O. Box 49, H-1525 Budapest 114

Hungary

E-mail: suto@szfki.hu

Abstract

We study the ground state $\Psi_0 = \sum a_X |X\rangle$ of N hard-core bosons on a finite lattice in configuration space, $X = \{x_1, \dots, x_N\}$. All a_X being positive, the ratios $a_X / \sum a_Y$ can be interpreted as probabilities $P_a(X)$. Let E_0 denote the energy of the ground state and $|\partial X|$ the number of nearest-neighbour particle-hole pairs in the configuration X . We prove the concentration of P_a onto X with $|\partial X|$ in a $\sqrt{|E_0|}$ -neighbourhood of $|E_0|$, show that the average of a_X over configurations with $|\partial X| = n$ increases exponentially with n , discuss fluctuations about this average, derive upper and lower bounds on E_0 and give an argument for off-diagonal long-range order in the ground state.

PACS: 0530J, 0550, 7510J

Appeared in *J. Phys. A: Math. Gen.* **34** (2001) 37-55, *ibid.* **34** (2001) 6209

1 Introduction

Bosons on a lattice interacting via an infinite on-site repulsion (hard-core bosons) represent a system of double interest. This is the simplest example of an interacting Bose gas and, thus, the most promising candidate for a rigorous treatment of Bose-Einstein condensation of interacting particles. The model is also known to be equivalent to a system of $\frac{1}{2}$ spins (Matsubara and Matsuda 1956) coupled via the X and Y and possibly the Z components of neighbouring spins and exposed to an external magnetic field in the Z direction. Ordering of the planar component of the spins is equivalent to Bose-Einstein condensation or the appearance of an off-diagonal long-range order (ODLRO, Yang 1962) in the system of bosons. Apart from some exceptions, such as bounds on the density of the condensate (Tóth 1991) or the discussion of the model on complete graphs (Tóth 1990, Penrose 1991), the most interesting rigorous results were formulated in spin terminology and obtained by the use of a particular symmetry, reflexion positivity (Dyson *et al* 1978, Kennedy *et al* 1988, Kubo and Kishi 1988). This introduced limitations on the value of the external field (zero field) and the lattice type (essentially hypercubic lattices). Expressed in terms of a boson gas, ODLRO was shown only at half filling on hypercubic lattices: in the ground state in dimensions ≥ 2 , and for positive temperatures above two dimensions. The proof of ordering does not offer much insight into the structure of the state. The only case in which details are known is the mathematically nice but physically trivial example of the complete graph (Tóth 1990, Penrose 1991).

In this paper we provide informations concerning the ground state Ψ_0 , valid for very different lattices and arbitrary particle densities. The main result is the proof of a strong non-uniformity of Ψ_0 in configuration space. It appears mathematically in the form of a large-deviation principle and makes the ground state resemble a thermal Gibbs distribution of a classical gas on the same lattice. It also leads to an approximate expression for Ψ_0 and to an argument for Bose-Einstein condensation in the ground state in two and higher dimensions.

Let $\mathbf{L}$ be an infinite lattice which, for the sake of simplicity, will be supposed to be regular with a constant coordination number (degree) k . Throughout the paper Λ denotes a finite connected part of $\mathbf{L}$ taken with periodic boundary conditions so as to keep the degree constant (not really essential). The Hamiltonian we are going to study in detail is

$$H_0 = - \sum_{\langle xy \rangle} (b_x^* b_y + b_y^* b_x) . \quad (1)$$

We write $x, y, \dots$ for the vertices (sites) and $\langle xy \rangle$ for the edges (nearest-neighbour pairs); b_x^* and b_x create, respectively, annihilate a hard-core boson at x . Boson operators at different sites commute with each other while

$$b_x^* b_x + b_x b_x^* = 1 \quad (2)$$

accounts for the hard-core condition. Correspondence with spin models is obtained by setting $b_x = S_x^-$ and $b_x^* = S_x^+$. The Hamiltonian conserves the number of bosons,

$$N = \sum_{x \in \Lambda} n_x = \sum_{x \in \Lambda} b_x^* b_x \quad (3)$$

and is also invariant under particle-hole transformation. We can, therefore, fix N so that $\rho = N/|\Lambda|$ is between 0 and $\frac{1}{2}$. (Here and below, if A is a finite set, $|A|$ stands for the number of elements.) Let $X, Y, \dots$ denote N -point subsets of Λ , also called configurations. A convenient basis is formed by the states

$$|X\rangle = \left(\prod_{x \in X} b_x^* \right) |0\rangle \quad (4)$$

where $|0\rangle$ is the vacuum state. According to the Perron-Frobenius theorem, there is a single ground state Ψ_0 whose all coefficients in the basis (4) are positive. Let $a = (a_X)$ denote the vector of these coefficients. Since the unique source of energy is hopping constrained by an on-site exclusion, we can expect that a_X increases, at least ‘on average’, with an increasing number of nearest neighbour particle-hole pairs. More

precisely, let ∂X denote the set of nearest-neighbour particle-hole pairs for X (the ‘boundary’ of X) and Ω_n the ensemble of configurations X with ‘boundary length’ $|\partial X| = n$. It is easily seen that

$$|\partial X| = \langle X | H_0^2 | X \rangle = \langle X | \Delta | X \rangle \quad (5)$$

where

$$\Delta = \sum_{\langle xy \rangle} [n_x(1 - n_y) + n_y(1 - n_x)] \quad (6)$$

is the diagonal part of H_0^2 , and $|\Omega_n|$ is the multiplicity of the eigenvalue n of Δ . We expect that the X -dependence of a_X appears mainly through $|\partial X|$, and the average

$$\alpha_n = \frac{1}{|\Omega_n|} \sum_{X \in \Omega_n} a_X \quad (7)$$

is some increasing function of n . We shall indeed find that α_n rapidly grows with n – exponentially fast for n of the order of N and even faster for smaller n . To make a comparison with the free Bose gas, let us note that the ground state of the latter reads

$$\Psi_{\text{free}} = |\Lambda|^{-N/2} \sum_{\{m_x \geq 0\}_{x \in \Lambda} : \sum m_x = N} \sqrt{\frac{N!}{\prod_{x \in \Lambda} m_x!}} \prod_{x \in \Lambda} (a_x^*)^{m_x} |0\rangle \quad (8)$$

where a_x^* is the ordinary boson creation operator at x . If we project out the part with no particle encounters, we find $\text{constant} \times \sum |X\rangle$, the sum going over the basis (4). This uniform sum is qualitatively very different from what we actually find for the ground state, and the energy of this state will also be found to be extensively higher than that of the ground state.

We start, in section 2, by studying the probability measure

$$p_a(n) = \frac{|\Omega_n| \alpha_n}{\sum_m |\Omega_m| \alpha_m} . \quad (9)$$

We show that the mean value according to p_a is $|E_0|$, the modulus of the ground state energy, while the mean square deviation D_a^2 is of the order of N , so that $p_a(n)$ is concentrated onto integers with $|n - |E_0||$ of the order of $\sqrt{N}$. In section 3 we evoke a large deviation principle for the Ising model on the same lattice, having the approximate form

$$q(n) \equiv \frac{|\Omega_n|}{\sum_m |\Omega_m|} \propto e^{-\frac{(n-M)^2}{2D^2}} . \quad (10)$$

Here $M = \overline{|\partial X|}$, the (arithmetic) mean of boundary lengths among N -point configurations (and also the modulus of the energy of the projected free-boson ground state) and D^2 is the corresponding mean square deviation of $|\partial X|$. We derive a formula for D^2 as a function of ρ and k ($M = k\rho(1 - \rho)|\Lambda|$ can be obtained trivially). In section 4 we use the results on p_a and q to give an approximate expression for α_n in terms of E_0 , D_a , M and D . We present an argument for the monotonicity of α_n and deduce from it upper and lower bounds on D_a/D , predicting $\alpha_n = \text{constant}$ if $|E_0|/M = 1$ asymptotically (as $|\Lambda|$ and N go to infinity). The variational treatment in section 5 shows that $|E_0|/M > 1$ in the thermodynamic limit. A notable exception, with $|E_0| = M = N(|\Lambda| - N)$, is the complete graph where by permutational symmetry we actually have $a_X = \text{constant}$. In section 5 we also present lower bounds on E_0 for all densities. The physical consequences of our findings are resumed in section 6. Here we provide an argument for the existence of ODLRO in the two and higher dimensional models. It is based on a hypothesis which also permits to estimate the dimension dependence of the deviations of a_X from the average $\alpha_{|\partial X|}$ and to conclude that these deviations are irrelevant above two dimensions. In section 7 we briefly discuss extensions to Hamiltonians with more complicated interactions.

2 Non-uniformity of the ground state

The results of this section are valid for any connected, not necessarily regular lattice with all coordination numbers $\leq k$.

Let $A = (A_{XY})$ denote the matrix of $-H_0$ in the basis (4). Then A is the adjacency matrix of a graph G whose vertices are the configurations, and two configurations X and Y form an edge of G if and only if their symmetric difference is an edge of Λ : $X \setminus Y \cup Y \setminus X = \{x, y\} = \langle xy \rangle$. We denote the largest eigenvalue of A by λ_1 and the corresponding eigenvector by $a = (a_X)$. The ground state energy and wave function are related to them through $E_0 = -\lambda_1$ and $\Psi_0 = \sum a_X |X\rangle$. The coefficients of the ground state wave vector a being positive, $a_X / \sum a_Y$ can be interpreted as a probability $P_a(X)$ of X . As we shall see, $P_a(X)$ is very far from being a uniform distribution that one can find on complete graphs.

A well-known fact concerning adjacency matrices is that $(A^n)_{XY}$ provides the number of walks of length n in G between X and Y (Biggs 1974). Let

$$W_n(X) = \sum_Y (A^n)_{XY} , \quad (11)$$

the number of walks of length n starting from X . First, we note that the expectation value of this number with respect to P_a ,

$$\langle W_n \rangle_a \equiv \sum_X W_n(X) a_X / \sum_X a_X = \lambda_1^n = \langle W_1 \rangle_a^n . \quad (12)$$

This follows by taking the X component of the vector equation $A^n a = \lambda_1^n a$, summing over X and dividing by $\sum a_X$. The above equality holds true for any simple graph: the probability measure P_a ‘sees’ the graph as if it were regular with a degree λ_1 .

Next, we make use of the fact that actually we are dealing with a sequence of graphs, $G = G_{\Lambda, N}$, having the particular property that the typical degree is of order N while its change between neighbouring vertices is of order one: The degree of X is its number of neighbours, $W_1(X)$. There is a one-to-one correspondence between nearest neighbour particle-hole pairs if X is considered as a subset of Λ and neighbours of X as a vertex in G . Therefore, ∂X can be identified with the set of neighbours of X in G and we have $W_1(X) = |\partial X|$. When passing from X to a $Y \in \partial X$ a neighbouring particle-hole pair is interchanged. For both the particle and the hole the number of neighbours of the opposite kind can change by at most $k - 1$, whence

$$|W_1(X) - W_1(Y)| \leq 2(k - 1) \quad \text{if } Y \in \partial X . \quad (13)$$

Now

$$W_2(X) = \sum_{Y \in \partial X} |\partial Y| \quad (14)$$

which together with (13) yields

$$|W_1(X)^2 - W_2(X)| \leq 2(k - 1)|\partial X| . \quad (15)$$

Taking the expectation value and using (12),

$$0 \leq D_a^2 \equiv \langle W_1^2 \rangle_a - \langle W_1 \rangle_a^2 \leq 2(k - 1)\lambda_1 . \quad (16)$$

By Chebyshev’s inequality we then find that for any $\varepsilon > 0$

$$P_a \left(|W_1 - \lambda_1| > \sqrt{2(k - 1)\lambda_1/\varepsilon} \right) < \varepsilon \quad (17)$$

or equivalently

$$\sum_{|W_1(X) - \lambda_1| \leq \sqrt{2(k - 1)\lambda_1/\varepsilon}} a_X \geq (1 - \varepsilon) \sum a_X . \quad (18)$$

Similarly to (14),

$$W_n(X) = \sum_{Y_1 \in \partial X} \sum_{Y_2 \in \partial Y_1} \cdots \sum_{Y_{n-1} \in \partial Y_{n-2}} |\partial Y_{n-1}| \quad (19)$$

which, together with (13), yields

$$W_n < \prod_{l=0}^{n-1} (W_1 + 2(k-1)l) \quad W_n > \prod_{l=0}^{n-1} (W_1 - 2(k-1)l) . \quad (20)$$

These bounds are nontrivial if $n \ll |\Lambda|$, for example, for n fixed and $|\Lambda|$ going to infinity. (In the opposite limit we have the stronger inequality $W_n(X) \leq (kN)^n$.) From (20) we get

$$|W_1^n - W_n| \leq \sum_{m=1}^{n-1} (2k-2)^m \left(\sum_{1 \leq l_1 < \cdots < l_m \leq n-1} l_1 \cdots l_m \right) W_1^{n-m} . \quad (21)$$

Taking the expectation value,

$$|\langle W_1^n \rangle_a - \lambda_1^n| \leq \sum_{m=1}^{n-1} (2k-2)^m \left(\sum_{1 \leq l_1 < \cdots < l_m \leq n-1} l_1 \cdots l_m \right) \langle W_1^{n-m} \rangle_a . \quad (22)$$

Replacing n by $n-m$ we obtain

$$\langle W_1^{n-m} \rangle_a \leq \lambda_1^{n-m} + O(\lambda_1^{n-m-1}) . \quad (23)$$

Now λ_1 is of order $|\Lambda|$, so finally

$$|\langle W_1^n \rangle_a - \lambda_1^n| \leq n(n-1)(k-1)[1 + O(|\Lambda|^{-1})]\lambda_1^{n-1} . \quad (24)$$

Equations (16)-(18) and (24) describe the concentration of the probability measure P_a on configurations X whose boundary length $|\partial X|$ is in a $\sqrt{N}$ -neighbourhood of $|E_0|$. We shall refer to this property as the non-uniformity of the ground state. Summing over X in Ω_n , P_a gives rise to the probability distribution p_a , which is, hence, peaked about $|E_0|$. In section 4 we shall arrive at the same conclusion in a different way, by studying first the *a priori* distribution $q(n) = |\Omega_n| / \sum_m |\Omega_m|$.

3 Large-deviation principle for the a priori distribution

The precise computation of $|\Omega_n|$ is a difficult and unsolved combinatorial problem. The logarithm of this number is the entropy of the Ising model, for energy n , in a microcanonical ensemble with a fixed magnetization, $\sum_{x \in \Lambda} \sigma_x = 2N - |\Lambda|$. Indeed, if we put $\sigma_x = 1$ for x in X and $\sigma_x = -1$ elsewhere, we get an Ising configuration with a density ρ of $+$ spins. The corresponding union of contours can be identified with ∂X whose total length $|\partial X|$ is the energy of the Ising configuration. For such systems there exists a strong version of the equivalence of ensembles which, applied to our case, states that the distribution q of $|\partial X|$ satisfies a large deviation principle whose rate function is, apart from a shift, minus the specific entropy $s(e, \rho)$ of the Ising model ($e = n/|\Lambda|$), cf. Pfister 1991, Dobrushin and Shlosman 1994 and Lewis *et al* 1994. The exact form of the entropy is unknown. To circumvent this problem, we use an approximate formula for the probability of having a boundary of length n ,

$$q(n) \approx Z^{-1} \exp\left\{-\frac{(n-M)^2}{2D^2}\right\} . \quad (25)$$

Here $Z \propto \sqrt{N}$ is for normalization and

$$M = \overline{|\partial X|} = k\rho(1-\rho)|\Lambda| \quad D^2 = \overline{(|\partial X| - \overline{|\partial X|})^2} , \quad (26)$$

cf. Eq. (10) in the Introduction. The Gaussian approximation is correct in a neighbourhood of the maximum of the specific entropy if $|\Lambda|/D^2$ is non-vanishing in the thermodynamic limit. This is what we are going to check by explicitly computing the mean square deviation of $|\partial X|$. We show that

$$D^2 = [k - 2(2k - 1)\rho(1 - \rho)]k\rho(1 - \rho)|\Lambda| = [k - 2(2k - 1)\rho(1 - \rho)]M . \quad (27)$$

We have found this expression first for d -dimensional hypercubic lattices, and have also checked it for the triangular, honeycomb and Kagomé lattices. The apparent independence of the details of the lattice is somewhat surprising because our derivation below needs knowledge of the rather different local neighbourhoods up to next-nearest neighbours. The only common feature of all these lattices seems to be that all sites are symmetry-related and thus equivalent. This alone should therefore suffice to prove Eq. (27). Let us note that, with the microcanonical temperature defined as $(\partial s/\partial e)^{-1}$, the maximum of the entropy corresponds to infinite temperature. Therefore, in a neighbourhood of the maximum we are at a safe distance to the ferro- and antiferromagnetic phase transitions, which could weaken our approximation through a breakdown (in finite volumes) of the concavity of the microcanonical entropy (Pleimling and Hüller 2000).

Equation (27) is obtained by filling the sites of Λ independently and with equal probability ρ . We expect smaller order corrections to appear if the computation is done with N fixed, $N/|\Lambda| = \rho$ (cf. Eq. (52) below). So in this section X is a random subset of Λ whose probability to be selected is $\rho^{|\Lambda|} (1 - \rho)^{|\Lambda| - |X|}$, and $n_x = n_x(X)$ is a random variable taking the value 1 if x is in X and 0 otherwise. Then all n_x are independent and take 1 with probability ρ and 0 with probability $1 - \rho$.

We define

$$f_x = n_x \sum_{y \in \partial x} (1 - n_y) \quad (28)$$

where ∂x denotes the set of neighbours of x in Λ . The boundary length of X is obtained (cf. Eqs. (5) and (6)) as

$$|\partial X| = \sum_{x \in \Lambda} f_x(X) . \quad (29)$$

Thus the mean value of (29) is

$$M = \sum_{x \in \Lambda} \overline{f_x} = k\rho(1 - \rho)|\Lambda| \quad (30)$$

as claimed earlier.

Let $d(x, y)$ denote the graph distance of x and y in Λ , i.e. the length of the shortest walk between them. Since f_x and f_y are independent if $d(x, y) > 2$, we find

$$D^2 = \sum_{x, y} \overline{(f_x - \overline{f_x})(f_y - \overline{f_y})} = \sum_{x, y: d(x, y) \leq 2} r(x, y) \quad (31)$$

$$r(x, y) = \overline{f_x f_y} - \overline{f_x}^2 . \quad (32)$$

The computation of the different terms is straightforward by observing that $n_x^2 = n_x$. The contribution of the diagonal terms $x = y$ is the same for any k -regular lattice. Namely,

$$\overline{f_x^2} = k\rho(1 - \rho) + k(k - 1)\rho(1 - \rho)^2 \quad (33)$$

$$\sum_{x \in \Lambda} r(x, x) = |\Lambda| r(x, x) = [k - (2k - 1)\rho + k\rho^2]M . \quad (34)$$

The contribution of nearest neighbour pairs depends on the number of triangles containing a given edge. If there are ℓ such triangles then

$$\overline{f_x f_y} = \rho^2 [\ell(1 - \rho) + [(k - 1)^2 - \ell](1 - \rho)^2] \quad (35)$$

$$\sum_{x,y: d(x,y)=1} r(x,y) = k|\Lambda| r(x,y) = \rho[\ell\rho - (2k-1)(1-\rho)]M . \quad (36)$$

If x and y are next-nearest neighbours to each other, they may have m common nearest neighbours. Then

$$\begin{aligned} \overline{f_x f_y} &= \rho^2[m(1-\rho) + (k^2 - m)(1-\rho)^2] \\ r(x,y) &= m\rho^3(1-\rho) . \end{aligned} \quad (37)$$

In d -dimensional hypercubic lattices ($k = 2d$) there are next-nearest neighbour pairs with $m = 1$ and $m = 2$. Their contribution to D^2 is

$$\begin{aligned} \sum_{x,y: d(x,y)=2} r(x,y) &= k|\Lambda| r(x,y)_{m=1} + 4\binom{d}{2}|\Lambda| r(x,y)_{m=2} \\ &= \rho^2 M + (k-2)\rho^2 M = (k-1)\rho^2 M . \end{aligned} \quad (38)$$

For the triangular lattice ($k = 6$)

$$\sum_{x,y: d(x,y)=2} r(x,y) = k|\Lambda| [r(x,y)_{m=1} + r(x,y)_{m=2}] = (k-3)\rho^2 M . \quad (39)$$

In the honeycomb lattice ($k = 3$) each site has six next-nearest neighbours, all of the type $m = 1$. So

$$\sum_{x,y: d(x,y)=2} r(x,y) = 2k|\Lambda| r(x,y)_{m=1} = (k-1)\rho^2 M . \quad (40)$$

In the Kagomé lattice ($k = 4$) there are eight next-nearest neighbours with $m = 1$:

$$\sum_{x,y: d(x,y)=2} r(x,y) = 2k|\Lambda| r(x,y)_{m=1} = (k-2)\rho^2 M . \quad (41)$$

Finally, we obtain D^2 by adding (34) and (36) with $\ell = 0$ and (38) for hypercubic lattices, (34) and (36) with $\ell = 2$ and (39) for the triangular lattice, (34) and (36) with $\ell = 0$ and (40) for the honeycomb lattice and (34), (36) with $\ell = 1$ and (41) for the Kagomé lattice. All yield (27).

If the random variables f_x were independent, the mean square deviation of their sum would be given by (34). For any $\rho \leq \frac{1}{2}$ this is larger than the actual value (27), so on average the f_x are negatively correlated. Because $\ell \leq k-1$, the nearest neighbour correlation $r(x,y) < 0$ for all $\rho \leq \frac{1}{2}$, cf. Eq. (36). On the other hand, according to (37), for next-nearest neighbours $r(x,y)$ is always positive. Apparently, the induced probability distribution for the f_x could be approximated by a Boltzmann-Gibbs distribution written with an antiferromagnetic nearest-neighbour Hamiltonian for the centralized variables $f_x - \overline{f_x}$ and with a temperature and an external field chosen so as to fit the computed average (30) and nearest-neighbour correlations (36). If the effective β and field are below their respective critical values we get exponentially decaying antiferromagnetic correlations, and the approximation is qualitatively correct. Adding next-nearest-neighbour interactions one could fit the computed next-nearest-neighbour correlations, and so on.

4 The form of the averaged wave vector

Using the large-deviation result for $q(n)$ we can obtain more precise information on $p_a(n)$ and α_n . Apart from normalization $p_a(n)$ is obtained by multiplying $q(n)$ with α_n . Since, as we shall see in the next section, $\lambda_1 > M$ and the difference is of the order of N , α_n has to increase exponentially fast – at least in a neighbourhood of M – so as to shift the expectation value M of q to the expectation value λ_1 of p_a . As a result, we obtain the approximate expression $p_a(n) \sim \exp -(n - \lambda_1)^2 / 2D_a^2$, consistent with our

findings in section 2. This is a second, independent, argument for the non-uniformity of the ground state, assuming the form of another large-deviation principle for $|\partial X|$. To get α_n we equate this form of $p_a(n)$ with the other one, (9), where $|\Omega_n|$ is estimated through (10). This gives

$$\alpha_n \sim \exp \left(\frac{(n-M)^2}{2D^2} - \frac{(n-\lambda_1)^2}{2D_a^2} \right) \propto \exp \frac{(D_a^2 - D^2)n^2 + 2(D^2\lambda_1 - D_a^2M)n}{2D^2D_a^2} \quad (42)$$

For the moment, we have only trivial estimates for λ_1 ($\geq M$) and D_a (cf. Eq. (16)). More precise bounds on D_a can be obtained from the monotonic increase of α_n .

The argument telling that α_n should increase for all allowed n is as follows. From the spectral decomposition of A^m we obtain (supposing $\sum a_X^2 = 1$)

$$a_X a_Y = \lim_{m \rightarrow \infty} \lambda_1^{-m} (A^m)_{XY} \quad (43)$$

if Λ is non-bipartite, and

$$a_X a_Y = \lim_{m \rightarrow \infty} \frac{1}{2} [\lambda_1^{-m} (A^m)_{XY} + \lambda_1^{-m-1} (A^{m+1})_{XY}] = \lim_{m \rightarrow \infty} \frac{\lambda_1^{-r_m}}{2} (A^{r_m})_{XY} \quad (44)$$

if Λ is bipartite; $r_m = 2m$ or $2m+1$ if the graph distance $d_G(X, Y)$ of X and Y is even or odd, respectively. Summing over Y and averaging over X in Ω_n we obtain, for example, for a non-bipartite Λ ,

$$\alpha_n = \left(\sum a_Y \right)^{-1} \lim_{m \rightarrow \infty} \lambda_1^{-m} \frac{1}{|\Omega_n|} \sum_{X \in \Omega_n} W_m(X) . \quad (45)$$

Inspecting Eq. (19) one can conclude that for any m the average of $W_m(X)$ over Ω_n increases with n and, hence, $\alpha_n \geq \alpha_{n'}$ if $n > n'$.

Let $n_{\min}$ and $n_{\max}$ denote the smallest and largest allowed values of $|\partial X|$, respectively. Since $n_{\min} = o(N)$, monotonicity of α_n implies through (42)

$$\frac{n_{\max} - \lambda_1}{n_{\max} - M} \leq \frac{D_a^2}{D^2} \leq \frac{\lambda_1}{M} \quad (46)$$

up to an error of $o(1)$. Therefore, the coefficient of n in the exponent of α_n is non-negative. Clearly $\lambda_1/M = 1$ would imply $D_a/D = 1$ and $\alpha_n = \text{constant}$. In the next section we show that for finite dimensional lattices $\lambda_1/M > 1$ asymptotically and, hence, the two bounds in (46) form an interval around 1. On bipartite lattices $n_{\max} = kN$, so at half filling $n_{\max} = 2M$ and 1 is in the centre of the interval. It is possible that $D_a = D$ and the exponent of α_n is linear in n . However, even if $D_a \neq D$, the quadratic term in the exponent is of the same order as or even smaller (if $n = o(N)$) than the linear term.

More information can be extracted from Eqs. (43) and (44) if we combine them with (18). If m is large enough,

$$1 - 2\varepsilon \leq \frac{1}{W_m(X)} \sum_{||\partial Y| - \lambda_1| \leq \sqrt{2(k-1)\lambda_1/\varepsilon}} (A^m)_{XY} \leq 1 , \quad (47)$$

that is, for any X and any sufficiently large m an overwhelming majority of walks of length m starting from X on the graph G end up in a vertex whose degree is $\lambda_1 + O(\sqrt{N})$. In the expression

$$W_m(X) = (a_X \sum a_Y) \lambda_1^m + o(\lambda_1^m) \quad (m \rightarrow \infty) , \quad (48)$$

obtained from (43), λ_1^m accounts for this ‘long-time’ behaviour. Let us note that (47) cannot be understood by imagining the walks on G as realizations of some simple random process. Because G is nonregular (unless Λ is a complete graph), no Markov process can assign equal probabilities to all walks of equal length. As an example, for a locally unbiased random motion (which chooses among neighbours

with equal probability) the most probable individual walks of a given length are the ‘descending’ ones, those going towards vertices of a lower degree, and the less probable the ‘ascending’ ones. Because of the form (25) of the distribution of degrees, an X with $|\partial X| < M$ has typically more ascending than descending neighbours and *vice versa* for $|\partial X| > M$. Therefore, the most probable degrees of end-points of very long walks would be nevertheless close to M (but not to λ_1).

Although the monotonicity of α_n holds for all n between $n_{\min}$ and $n_{\max}$, the validity of (42) is limited to a neighbourhood of the interval $[M, \lambda_1]$. In particular, the growth of α_n for $n = o(N)$ is much steeper than the exponential predicted by (42). Indeed, let us write the eigenvalue equation in the form

$$\frac{1}{|\partial X|} \sum_{Y \in \partial X} a_Y = \frac{\lambda_1}{|\partial X|} a_X . \quad (49)$$

For a d -dimensional lattice $n_{\min} = O(N^{\frac{d-1}{d}})$. If we choose $|\partial X| = n$ to be of this order, we find that the average of a_Y over ∂X yields $N^{1/d}$ times a_X ! Since $q(n)$ is also rapidly increasing here, the average over ∂X is dominated by Y with $n+2 \leq |\partial Y| \leq n+2k-2$. From this and the monotonicity of α_n we conclude that α_{n+2k-2}/α_n is at least of order $N^{1/d}$.

5 Bounds on the ground state energy

5.1 Variational upper bounds

Variational estimates of the ground state energy are of the form

$$E_0 \leq \langle \psi | H | \psi \rangle / \langle \psi | \psi \rangle . \quad (50)$$

A trivial choice is

$$\psi = \sum |X\rangle \quad (51)$$

with the summation going over all N -point subsets of Λ . It yields $E_0 \leq -M$ where

$$M = \overline{|\partial X|} = \frac{2|EG|}{|VG|} = \frac{2|E\Lambda| \binom{|\Lambda|-2}{N-1}}{\binom{|\Lambda|}{N}} = k(1-\rho)N + O(1) , \quad (52)$$

cf. Eq. (30). Here $|EG|$ and $|VG|$ denote the number of edges and vertices of G , respectively, and $|E\Lambda|$ the number of edges of Λ . It is not *a priori* obvious that this bound can be improved in the order of the volume, and it is important to know that it really can. In the opposite case, if $-k\rho(1-\rho)$ were the true ground state energy per site then, as in the complete graph, the Hamiltonian (1) would have a product ground state in an infinite volume,

$$\Psi = \prod_x (\sqrt{\rho} |n_x = 1\rangle + \sqrt{1-\rho} |n_x = 0\rangle) , \quad (53)$$

i.e. no local perturbation could decrease the energy of Ψ . (Because of the product structure, if the energy could be decreased locally, the specific energy $-k\rho(1-\rho)$ of Ψ could also be decreased.) Since Ψ shows ODLRO with the value of the order parameter at its theoretical maximum ($\rho(1-\rho)$, cf. Appendix), by proving it cannot be a ground state we exclude a trivial scenario for Bose-Einstein condensation.

To prove that $|E_0| - M$ is of order N we apply trial functions of the form

$$\psi_v = \sum v_X |X\rangle \quad v_X = v(|\partial X|) \quad (54)$$

i.e. v_X depending on $|\partial X|$ only. The variational bound (50) reads $\lambda_1 \geq B(v)$ where

$$B(v) \equiv \frac{(v, Av)}{(v, v)} = \frac{\sum_n v(n) \sum_{i=-k+1}^{k-1} v(n+2i) \sum_{X \in \Omega_n} N_i(X)}{\sum_n v(n)^2 |\Omega_n|} . \quad (55)$$

Here $N_i(X)$ is the number of those neighbours of X having a boundary length $|\partial X| + 2i$. In Eq. (55) we have used (13) and the fact that the parity of $|\partial X|$ is the same for all X : even if N is even and that of k if N is odd.

The form (42) of α_n suggests that the best choice for $v(n)$ would be e^{xn+yn^2} with x of order 1 and y of order $\frac{1}{N}$. However, we do not expect the quadratic term to yield a significant improvement, and choose a simple exponential $v(n) = e^{xn}$ with $0 < x < x_{\max}$ where

$$x_{\max} = \frac{n_{\max} - M}{2D^2} . \quad (56)$$

Then by making use of the large-deviation principle for $|\partial X|$ we find

$$B(v) = \langle n \rangle_x \langle e^{2xi} \rangle_{[\langle n \rangle_x]} \quad (57)$$

where

$$\langle n \rangle_x = \frac{\sum n e^{2xn} |\Omega_n|}{\sum e^{2xn} |\Omega_n|} \quad \langle f(i) \rangle_n = \sum_{i=-k+1}^{k-1} f(i) \frac{1}{n |\Omega_n|} \sum_{X \in \Omega_n} N_i(X) . \quad (58)$$

Now, since $\langle n \rangle_0 = M$ and $d\langle n \rangle_x / dx \big|_{x=0} = 2D^2$, we find

$$\langle n \rangle_x = M(1 + 2xD^2/M + O(x^2)) . \quad (59)$$

We note that the approximate form (10) would yield the same result.

Next, we turn to the second term of $B(v)$. Obviously $\langle e^{2xi} \rangle_n \geq e^{-2(k-1)x}$. This yields $B(v)/M > 1$ for ρ near 0 but not near $\frac{1}{2}$. However, we can use Jensen's inequality $\langle e^{2xi} \rangle_n \geq e^{2x\langle i \rangle_n}$ together with the fact that $\langle i \rangle_{[M]}$ goes to zero as the volume increases. We then conclude that asymptotically $\langle i \rangle_{[\langle n \rangle_x]} = -\text{constant} \times x + \text{higher order terms}$, so that the average of the exponential in Eq. (57) equals $1 - O(x^2)$. Thus, for x small enough we indeed find $B(v)/M > 1$.

With somewhat more effort one can actually compute a lower bound on $B(v)$. We do not make it here, only notice that in optimizing such a bound according to x (and also in optimizing a lower bound on E_0 , see below) the knowledge of $n_{\max}$ is necessary. On bipartite lattices $n_{\max} = k\rho|\Lambda|$ for all $\rho \leq \frac{1}{2}$, and it is an easy graphical exercise to see that the same equality holds for $\rho \leq \frac{1}{3}$ on the triangular and Kagomé lattices. However, for both lattices $n_{\max}$ is constant between the densities $\frac{1}{3}$ and $\frac{1}{2}$: $2|\Lambda|$ for the triangular and $\frac{4}{3}|\Lambda|$ for the Kagomé lattice. This can be seen from the following argument. In general, $-n_{\max}$ is the ground state energy of the *antiferromagnetic* Ising model under the restriction that the magnetization is fixed, $\sum_{x \in \Lambda} \sigma_x = (2\rho - 1)|\Lambda|$. However, we do not need to deal with the restriction. In both cases the (unrestricted) ground state is known to be highly degenerate. Among the exponentially large number of ground state configurations there are non-magnetized ones, corresponding to $\rho = \frac{1}{2}$, others with a concentration of up-spins $\rho = \frac{1}{3}$, and between these two limits ρ can vary by steps of $1/|\Lambda|$. The rule is to flip zero-energy spins one by one. The common energy of all these configurations is easy to compute from the fact that in each triangle there is precisely one unsatisfied bond. This fixes the value of $n_{\max}$ as given above.

We also note that the optimal x (which maximizes $B(v)$) is small: of order 0.1 or smaller. We shall evoke this fact in the discussion of section 6.

5.2 Lower bounds

In their paper Dyson, Lieb and Simon (DLS) gave a lower bound on the ground state energy of the spin- $\frac{1}{2}$ XY-model (Dyson *et al* 1978, Theorem C.1). Because it corresponds to an upper bound on the norm of the Hamiltonian (1) in Fock space, it is automatically valid for the hard-core boson gas (1) at any density. It reads

$$|E_0| \leq \frac{1}{4}|\Lambda| \begin{cases} \sqrt{k(k+2)} & \text{if } k \text{ is even} \\ k+1 & \text{if } k \text{ is odd} . \end{cases} \quad (60)$$

The above bound is to be compared with the trivial bound $|E_0| \leq n_{\max}$ obtained by writing Eq. (49) for an $X = X_0$ which maximizes a_X . We then see that (60) is nontrivial for the d -dimensional hypercubic lattice if $\rho > \frac{1}{4}\sqrt{1 + \frac{1}{d}}$, for the triangular lattice if $\rho > \frac{1}{2\sqrt{3}}$, for the honeycomb lattice if $\rho > \frac{1}{3}$ and for the Kagomé lattice if $\rho > \frac{1}{4}\sqrt{\frac{3}{2}}$.

The bound (60) is the best at $\rho = \frac{1}{2}$ and can be improved for lower densities.

If e_0 is the ground state energy per site then

$$|e_0| = \langle \Psi_0 | A_x | \Psi_0 \rangle \equiv \langle \Psi_0 | \frac{1}{2} \sum_{y \in \partial x} (b_x^* b_y + b_y^* b_x) | \Psi_0 \rangle \quad (61)$$

for any x in Λ . A_x preserves the number of bosons in $\Lambda_x = \{x\} \cup \partial x$, therefore it commutes with $P_{x,j}$, the orthogonal projection to the subspace $\mathcal{H}_{x,j}$ of states with j particles in Λ_x , where $j = 0, 1, \dots, k+1$. We can write

$$\langle \Psi_0 | A_x | \Psi_0 \rangle = \sum_{j=0}^{k+1} \frac{\langle \Psi_0 | P_{x,j} A_x P_{x,j} | \Psi_0 \rangle}{\langle \Psi_0 | P_{x,j} | \Psi_0 \rangle} \langle \Psi_0 | P_{x,j} | \Psi_0 \rangle . \quad (62)$$

Since $\langle \Psi_0 | P_{x,j} \cdot P_{x,j} | \Psi_0 \rangle / \langle \Psi_0 | P_{x,j} | \Psi_0 \rangle$ is a normalized positive linear functional,

$$\langle \Psi_0 | A_x | \Psi_0 \rangle \leq \sum_j \lambda_{\max}(j) \langle \Psi_0 | P_{x,j} | \Psi_0 \rangle , \quad (63)$$

where $\lambda_{\max}(j)$ is the maximum eigenvalue of A_x restricted to $\mathcal{H}_{x,j}$. Now $\lambda_{\max}(j)$ was computed by DLS:

$$\lambda_{\max}(j) = \frac{1}{2} \sqrt{j(k+1-j)} . \quad (64)$$

The DLS-bound (60) corresponds to the maximum of (64). Let $h(t)$ be obtained by linear interpolation through the points $(j, \lambda_{\max}(j))$:

$$h(t) = \frac{1}{2} \left\{ ([t] + 1 - t) \sqrt{[t](k+1-[t])} + (t - [t]) \sqrt{([t] + 1)(k - [t])} \right\} . \quad (65)$$

Since $h(t)$ is a concave function, and the right member of (63) can be written as $\int h(t) d\nu(t)$ with a probability measure ν concentrated on the integers from 0 to $k+1$, by Jensen's inequality

$$|e_0| \leq h \left(\int t d\nu(t) \right) = h \left(\sum_{j=0}^{k+1} j \langle \Psi_0 | P_{x,j} | \Psi_0 \rangle \right) . \quad (66)$$

The expectation value in the argument of h can be evaluated and yields $(k+1)\rho$. Indeed, using the translation invariance of Ψ_0 , after a simple algebra we find

$$\sum_{j=0}^{k+1} j \langle \Psi_0 | P_{x,j} | \Psi_0 \rangle = \sum_X a_X^2 \frac{1}{|\Lambda|} \sum_{x \in \Lambda} |X \cap \Lambda_x| . \quad (67)$$

Let χ_X and χ_{Λ_x} denote the characteristic functions of X and Λ_x , respectively. Then

$$|X \cap \Lambda_x| = \sum_{y \in \Lambda} \chi_X(y) \chi_{\Lambda_x}(y) \quad (68)$$

and therefore

$$\sum_{x \in \Lambda} |X \cap \Lambda_x| = \sum_{y \in \Lambda} \chi_X(y) \sum_{x \in \Lambda} \chi_{\Lambda_x}(y) = |X| \cdot |\Lambda_x| = N(k+1) . \quad (69)$$




Figure 1: Approximations of the ground state energy per site as a function of the density in the square lattice.

Hence, we obtain

$$|e_0| \leq \min\{h((k+1)\rho), k\rho\} \leq \frac{1}{2}(k+1)\sqrt{\rho(1-\rho)} \quad (70)$$

where the last member results from majorizing $h(t)$ by $\frac{1}{2}\sqrt{t(k+1-t)}$. Taking the minimum in Eq. (70) is not superfluous. For $\rho < \frac{1}{k+1}$ we have $h((k+1)\rho) = \frac{1}{2}\sqrt{k}(k+1)\rho$. Since $n_{\max}/|\Lambda| = k\rho$ is smaller for all k , the trivial bound is better for small densities. Actually the trivial bound provides the right asymptotics at $\rho = 0$ because $e_0 = -k\rho + O(\rho^2)$ near $\rho = 0$, as one can see by comparing the trivial upper and lower bounds, $-M/|\Lambda|$ and $-n_{\max}/|\Lambda|$.

Numerical works on the spin- $\frac{1}{2}$ XY model in the square lattice yield an approximate formula,

$$e_0(\rho) = -1.09766 + 4.835(0.5 - \rho)^2 - 1.99232(0.5 - \rho)^4 + 0.85952(0.5 - \rho)^6. \quad (71)$$

The first two terms come respectively from a Monte Carlo (Zhang and Runge 1992) and a finite-size scaling (Hamer *et al* 1999) study, to which we have added the fourth and the sixth order terms to obtain $e_0(0) = 0$ and $e'_0(0) = -4$. In Figure 1 we have plotted the trivial upper bound $-4\rho(1-\rho)$, the formula (71) and the lower bound $-\min\{h((k+1)\rho), k\rho\}$. We have also shown numerical points for a 5×5 lattice from Table 1 of Hamer *et al* 1999. They nicely follow the curve (71).

Using the formula (71) we can compute the lower and upper bounds (46) on D_a^2/D^2 . Both of them increase monotonically as ρ varies between 0 and 0.5: the lower bound from ~ 0.66 to 0.90234 and the upper bound from 1 to 1.09766, their difference decreasing monotonically at the same time.

6 Off-diagonal order and the final form of the ground state

Off-diagonal long-range order in the ground state is characterized by (cf. Appendix)

$$\omega_0 \equiv \lim \frac{1}{|\Lambda|^2} \sum_X a_X \sum_{x \in X} \sum_{y \notin X} a_{X \cup \{y\} \setminus \{x\}} > 0 \quad (72)$$

where $\lim$ means the thermodynamic limit ($|\Lambda|$ and N going to infinity and $N/|\Lambda|$ going to ρ). In (72) $\sum a_X^2 = 1$ is supposed.

Let us see, what would we obtain if a_X did not fluctuate within Ω_n , i.e. if $a_X = \alpha_{|\partial X|}$. This and (42) would imply another large-deviation principle in the approximate form

$$\sum_{X \in \Omega_n} a_X^2 = \alpha_n^2 |\Omega_n| \sim \exp \left[-\frac{(n - \lambda_2)^2}{2D_{a^2}^2} \right] \quad (73)$$

with

$$\lambda_2 = \lambda_1 + D_a^2 \frac{\lambda_1 - M}{2D^2 - D_a^2} \quad D_{a^2}^2 = \frac{D^2 D_a^2}{2D^2 - D_a^2}. \quad (74)$$

The mean value λ_2 is larger than λ_1 because

$$\frac{D_a^2}{D^2} \leq \frac{\lambda_1}{M} < \min \left\{ \frac{1}{1 - \rho}, \frac{k + 1}{2k\sqrt{\rho(1 - \rho)}} \right\} < 2, \quad (75)$$

see Eq. (70). As

$$||\partial(X \cup \{y\} \setminus \{x\})| - |\partial X|| \leq 2k, \quad (76)$$

with $a_{X \cup \{y\} \setminus \{x\}} \geq \alpha_{|\partial X| - 2k}$ we would find

$$\omega_0 \geq \rho(1 - \rho) \lim e^{-\frac{2k(\lambda_1 - M)}{2D^2 - D_a^2}} > 0. \quad (77)$$

The argument above, together with the conclusion (77) may be right in high enough dimensions but it is known to be incorrect in one dimension. Fluctuations have to be taken into account.

A way to prove (72) would be to show

$$a_{X \cup \{y\} \setminus \{x\}} \geq c a_X \quad (78)$$

for all X , x and y with a $c > 0$ independent of Λ and N . If (78) did hold true, we would find $\omega_0 \geq c\rho(1 - \rho) > 0$. In the complete graph (78) is verified with $c = 1$ and equality sign. However, we know already from the considerations following Eq. (49) that in finite dimensional lattices (78) cannot hold indeed for all X : e.g., in the case when $|\partial X| \propto N^{1-\frac{1}{d}}$, (78) can be satisfied only with $c \propto N^{-\frac{1}{d}}$. Also, we should not be able to prove (78) in one dimension. Below we present an argument which takes into account fluctuations and distinguishes between one, two and higher dimensions.

First, we note that we need (78) only for X in a subset $S(\Lambda, N)$ such that the sum of a_X^2 over S is non-vanishing in the thermodynamic limit. Let us introduce a function $R(X)$ by setting $a_X = \alpha_{|\partial X|} e^{R(X)}$. Then

$$\sum_{X \in \Omega_n} e^{R(X)} = |\Omega_n| \leq \sum_{X \in \Omega_n} e^{2R(X)} \leq |\Omega_n|^2, \quad (79)$$

so we can define an ϵ_n such that $0 \leq \epsilon_n \leq 1$ and

$$\sum_{X \in \Omega_n} e^{2R(X)} = |\Omega_n|^{1+\epsilon_n}. \quad (80)$$

Using this and Eq. (42) we find

$$\sum_{X \in \Omega_n} a_X^2 = \alpha_n^2 |\Omega_n|^{1+\epsilon_n} \sim \exp \left[-\frac{(n - M_n)^2}{2D_n^2} \right] \quad (81)$$

with

$$M_n = \lambda_1 + \frac{(1 - \epsilon_n)D_a^2}{2D^2 - (1 - \epsilon_n)D_a^2}(\lambda_1 - M) \quad D_n^2 = \frac{D^2 D_a^2}{2D^2 - (1 - \epsilon_n)D_a^2} . \quad (82)$$

Because of the possible n -dependence of M_n and D_n , Eq. (81) may not describe a large deviation principle. However, M_n and D_n satisfy the inequalities $\lambda_1 \leq M_n \leq \lambda_2$ and $D_a/\sqrt{2} \leq D_n \leq D_{a^2}$ and, hence, for any $\varepsilon > 0$

$$\lim_{(1-\varepsilon)\lambda_1 \leq |\partial X| \leq (1+\varepsilon)\lambda_2} \sum a_X^2 = 1 . \quad (83)$$

For S we can take the set of summation with any $\varepsilon \geq 0$.

Next, we observe that by using Eqs. (43) and (44) we can write

$$\frac{a_X}{a_Y} = \lim_{m \rightarrow \infty} \frac{W_m(X)}{W_m(Y)} \quad \text{or} \quad \lim_{m \rightarrow \infty} \frac{\lambda_1 W_m(X) + W_{m+1}(X)}{\lambda_1 W_m(Y) + W_{m+1}(Y)} \quad (84)$$

for non-bipartite or bipartite lattices, respectively. The convergence in (84) is usually slow. However, we expect that if $|\partial X|$ and $|\partial Y|$ do not differ too much then, at least up to the order of magnitude, a_X/a_Y can be approximated by the ratio of the number of walks whose length is the distance between X and Y . Thus, we conjecture that for $|\partial X| = c_1 N$ and $||\partial Y| - |\partial X|| \leq c_2$

$$\left| \ln \frac{a_X}{a_Y} - \ln \frac{W_{d_G(X,Y)}(X)}{W_{d_G(X,Y)}(Y)} \right| \leq c_3 \quad (85)$$

where c_3 may depend on the constants c_1 and c_2 but not on the size of the system. In what follows, we examine the consequences of this hypothesis.

If the aspect ratios of Λ are kept bounded, its diameter L is of order $|\Lambda|^{\frac{1}{d}}$ and the diameter of G is of order L^{d+1} . Over such a distance (85) allows a gigantic change of a_X . However, the distance to be considered for (78) is much smaller. If the symmetric difference of X and Y is $\{x, y\}$, it is easily seen that $d_G(X, Y) = d(x, y) \leq L$. Let $|\partial X| = n \propto N$. Above one dimension we apply the inequalities (20) to obtain

$$\frac{W_{d_G(X,Y)}(X)}{W_{d_G(X,Y)}(Y)} \leq \left(\frac{1 + 2(k-1)L/n}{1 - [2(k-1)L + 2]/n} \right)^L . \quad (86)$$

In the thermodynamic limit the upper bound remains finite in two dimensions and converges to 1 in higher dimensions, so we get the necessary estimate (78). In one dimension there exists no convergent upper bound. Thus, we find ODLRO in the ground state in two and higher dimensions but not in one dimension.

The difference between 2 and $2 + \varepsilon$ dimensions in the convergence of the right-hand side of Eq. (86) indicates that $d = 2$ is the critical dimension above which the deviations of a_X from the average $\alpha_{|\partial X|}$ are irrelevant. This remark can be made quantitative if we observe an interesting consequence of the hypothesis (85) on $R(X)$. If $|\partial X| \propto N$ then inside a sphere of radius $\propto \sqrt{N}$, centred at X , for all Y such that $||\partial Y| - |\partial X||$ is bounded by some constant of the order of unity $|R(Y) - R(X)|$ also remains below another constant of order 1. Comparing the scale $\sim \sqrt{N}$ with the diameter $\sim N^{1+1/d}$ of G we conclude that

$$|R(X)| \leq \max R(Y) - \min R(Y) < \text{const} \times N^{\frac{1}{2} + \frac{1}{d}} . \quad (87)$$

The maximum and the minimum are taken over $\Omega_{|\partial X|}$. The first inequality holds because for any n the average of $R(X)$ over Ω_n is negative, but $R(X)$ cannot be negative for all X .

According to Equation (87),

$$a_X = \exp[F(|\partial X|) + R(X)] = \exp \left[F(|\partial X|) + O \left(N^{\frac{1}{2} + \frac{1}{d}} \right) \right]. \quad (88)$$

Therefore, in one dimension $R(X)$ can dominate $F(|\partial X|)$, in two dimensions it can vary on the same scale $\sim N$, and above two dimensions the variation of $R(X)$ is negligible compared with that of $F(|\partial X|)$. Apart from an additive constant necessary for normalization, the approximate form of $F(n) = \ln \alpha_n$ can be read off from Eq. (42).

The upper bound in the right-hand member of Eq. (87) could, of course, overestimate the order of magnitude of the maximum of $|R(X)|$. However, because it seems to correctly distinguish between one, two and higher dimensions, it is probably sharp. This implies also that on the scale $\sim \sqrt{N}$ the function $R(X)$ can be considered as describing a ballistic motion over G rather than a random fluctuation about a negative average, which would lead to a smaller bound. Accepting ballisticity on smaller scales as well, one can find the order of magnitude of local variations of $R(X)$. We distinguish between two cases.

(a) If Λ is bipartite with sublattices V_1, V_2 then G is also bipartite with sublattices $\Gamma_1 = \{X : |X \cap V_1| \text{ is even}\}$ and $\Gamma_2 = \{X : |X \cap V_1| \text{ is odd}\}$. Along each path in G the subsequent terms belong to alternating sublattices. If $\partial^2 X$ denotes the set of second neighbours of X then

$$|R(X) - R(Y)| < \text{const} \times N^{-1/2} \quad \text{if } Y \in \partial^2 X \text{ and } |\partial Y| = |\partial X|. \quad (89)$$

(b) If Λ is not bipartite, neither is G , and no systematic compensation of terms of different signs is possible. In this case

$$|R(X) - R(Y)| < \text{const} \times N^{-1/2} \quad \text{if } Y \in \partial X \text{ and } |\partial Y| = |\partial X|. \quad (90)$$

The result on ODLRO could be obtained from (89), (90) as well.

As the two-dimensional case is marginal, some subtle logarithmic corrections, that we do not see, may modify our conclusions.

Above two dimensions we also find, as a further consequence of Eq. (87), that ϵ_n identically vanishes in the thermodynamic limit. Thus, $M_n/\lambda_2 \rightarrow 1$ and $D_n/D_{a^2} \rightarrow 1$, Eq. (81) asymptotically coincides with Eq. (73), Eq. (77) provides a valid lower bound on the order parameter, and Eq. (83) can be replaced by

$$\lim_{(1-\varepsilon)\lambda_2 \leq |\partial X| \leq (1+\varepsilon)\lambda_2} \sum a_X^2 = 1 \quad (91)$$

for any $\varepsilon > 0$. This property makes a_X^2 similar to a thermal Gibbs state for a classical lattice gas. The classical Hamiltonian (with $1/k_B T$ incorporated) is $H_{\text{eff}}(X) = -2F(|\partial X|) - 2R(X)$, and there is asymptotic equivalence between the canonical distribution and a microcanonical one concentrated on X 's with $H_{\text{eff}}(X)$ in an $o(N)$ neighbourhood of the canonical expectation value of H_{eff} . This holds in spite of H_{eff} containing long-range interactions. Indeed, $-F$ is a sum of nearest neighbour repulsive two-body interactions, f_x , and, if $D_a \neq D$, properly normalized long-range four-body interactions, $f_x f_y$, see Eqs. (28), (29) and (42), the whole yielding a global repulsion which tends to maximize $|\partial X|$. Now R may also include up to N body interactions but because $R(X) = o(N)$ and because a_X^2 corresponds to a high-temperature Gibbs state, it can be neglected: The fact that R may partly split the ground state degeneracy of $-F$ is irrelevant at high temperatures.

We have no rigorous proof that the limiting effective Gibbs state is a pure (in particular, a high-temperature) state. For this we should show that for all densities and all relevant n , $2F(n)/n < K_c$, the critical coupling of the Ising model on the same lattice. Even if we knew λ_1 exactly, our bounds (46) on D_a/D are not good enough to obtain this information. For example, in the case of the square lattice we can use the numerical fit (71). If we assume that $D_a \geq D$, we get $2F(n)/n \leq 2(\lambda_1 - M)/D^2$ where the upper bound increases with ρ and its maximum at $\rho = 0.5$ is $0.3906 < K_c = 0.4407$. On the other hand, if we use the lower bound for D_a , at half filling we find for $n \geq \lambda_1$ that $2F(n)/n \leq 0.628 > K_c$. An indication to the high-temperature character can be found in the variational estimates, cf. the final

remark of section 5.1. In the light of Eq. (88), trial functions of the form (54) acquire a particular importance: For $d > 2$ one could find with them the exact ground state energy per site, and the optimal exponential ansatz may not be far from the true ground state. The variable x appearing in the formula (57) corresponds to $\frac{1}{2}\beta J$ of the Ising model. The estimates we have done predict a value well below the critical one in any dimension: $\beta J < 0.2$ for all densities at $k = 3$, and decreases with increasing k , reaching 0 in the complete graph.

Now the fact that a_X^2 is equivalent to a high-temperature Gibbs state means that there is no classical (diagonal) order in the ground state, coexisting with the purely quantum-mechanical (off-diagonal) one. Diagonal order is characterized by an order parameter operator O which is diagonal in the basis (4) and for which the ground state expectation value $|\Lambda|^{-2}\langle\Psi_0|O^*O|\Psi_0\rangle = |\Lambda|^{-2}\sum a_X^2|O(X)|^2$ has a non-vanishing thermodynamic limit. A typical example is $O_q = \sum_x e^{i(q,x)}n_x$, associated with a periodic order with a wave vector q . Due to the correspondence with a Gibbs state, diagonal order in the ground state of the Bose gas is equivalent to low-temperature order in the associated classical lattice gas. The ground state of $-F$ is exponentially degenerate for $\rho < 1/2$. However, it is only two-fold degenerate on bipartite lattices at $\rho = 1/2$, and the degeneracy is not split by the translation invariant $R(X)$. Apparently, there is no qualitative argument against a crystalline order at half filling in the ground state of the $d > 2$ dimensional purely hard-core Bose gas. The nearest neighbour repulsion in $-F$ is simply not strong enough, reflecting the fact that a purely on-site repulsion in H_0 , even though infinite, cannot induce such an order.

7 Extensions and concluding remarks

The nonuniformity of the ground state we have found in a purely hard-core gas seems to be a general property of finite-range Hamiltonians. The extension to more complicated hard-core Hamiltonians is immediate. Let

$$\begin{aligned} H &= -\sum_{\langle xy \rangle} (b_x^* b_y + b_y^* b_x) + \sum V_x n_x - J \sum_{\langle xy \rangle} \left(n_x - \frac{1}{2}\right) \left(n_y - \frac{1}{2}\right) + K(\{n_x\}) \\ &\equiv H_0 + V + \frac{J}{4}(2\Delta - |E\Lambda|) + K \end{aligned} \quad (92)$$

where Δ is the diagonal part of H_0^2 , cf. Eq. (6), and K may contain further interactions of a bounded range. Because the terms additional to H_0 are diagonal in the basis (4), the Perron-Frobenius theorem still applies and the ground state wave vector $a_X > 0$. If A is the matrix of $-H$ then keeping the definition (11) we still have (12), and the inequalities of section 2 also hold with suitable modifications. In particular, in (13) and (15) $2(k-1)$ is to be replaced by $\kappa = |2 - J|(k-1) + \delta V + \delta K$ where $\delta V = \max_{\langle xy \rangle} |V_x - V_y|$ and $\delta K = \max_{X, Y \in \partial X} |K(Y) - K(X)|$ are of the order of unity, and in (16)-(18) we have $\kappa \langle |\partial X| \rangle_a$ instead of $2(k-1)\lambda_1$. Now

$$W_1(X) = A_{XX} + |\partial X| = \left(1 - \frac{J}{2}\right) |\partial X| - V(X) - K(X) + \frac{J}{4} |E\Lambda| \quad (93)$$

may not be positive and $\langle W_1 \rangle_a = \lambda_1 = -E_0$ may not be of order N . Therefore, there may not be a large deviation principle for the pair (W_1, a) . However, if we replace H by $H^c = H + cI$ where I is the identity operator, the ground state vector a will not change. The corresponding W_1 is $W_1^c(X) = W_1(X) - c$, and $D_a^2(W_1^c)$ is also independent of c . Let us choose $c = \langle A_{XX} \rangle_a$, then

$$\lambda_1 + c \equiv \lambda_1^c = \langle W_1^c \rangle_a = \langle |\partial X| \rangle_a \quad (94)$$

and, thus, we have a large deviation principle for (W_1^c, a) in the same form as for the purely hard-core interaction.

What really counts for the non-uniformity of a is not the large deviation principle for (W_1, a) but the variation of $W_1(X)$ over the set of N -point configurations. The ground state is nonuniform if

$[\max W_1(X) - \min W_1(X)]/\sqrt{N}$ goes to infinity with N . As an example, let us consider the nearest neighbour anisotropic Heisenberg model ($V = K = 0$). At $J = 2$, when it is isotropic, for any fixed N we have $W_1(X) \equiv |E\Lambda|/2$ and hence $\lambda_1 = |E\Lambda|/2$ and a_X is constant. Away from the isotropy point ΔW_1 is of order N and a is nonuniform.

If $\Omega_n = \{X : W_1(X) = w_n\}$, where $w_1 < w_2 < \dots$ are the possible values of W_1 , one can write down an approximate formula similar to Eq. (10), namely $q(n) \sim \exp[-(w_n - \overline{W_1})^2/2D^2(W_1)](w_{n+1} - w_n)$. Interpreting α_n as the average of a_X over the newly defined Ω_n , we have also the analogue of Eq. (42).

As earlier, we can write $a_X = \exp[F(W_1(X)) + R(X)]$, and Eq. (79) remains valid. The approximate form of F is

$$F(W_1(X)) \approx \frac{(W_1(X) - \overline{W_1})^2}{2D(W_1)^2} - \frac{(W_1(X) + E_0)^2}{2D_a(W_1)^2} + \text{const} . \quad (95)$$

One can replace the infinite on-site repulsion by a finite one provided that stability is assured, $H > -\text{const} \times N$. The dimension of the N -particle subspace will increase to $\binom{|\Lambda|+N-1}{N}$. A configuration X becomes a list of N elements containing vertices of Λ with possible repetitions that we can write as $X = \{x^{n_x}\}_{x \in \Lambda}$. The neighbourhood relation remains the same: $Y \in \partial X$ if Y can be obtained from X by moving a single particle along an edge of Λ , say, from x into y . The corresponding matrix element of $-H$ is then $A_{XY} = \sqrt{n_x(X)n_y(Y)}$. Now $\sum_{Y \neq X} A_{XY}$ can vary between $k\sqrt{N}$ and $\sqrt{2}kN$ when $|\partial X|$ varies between k and kN . However, A_{XX} can be of order N^2 and its change between neighbours of order N . Therefore, we miss a pointwise bound like (13) and cannot repeat the argument of section 2. Nevertheless, $-W_1(X) = \sum_Y \langle Y | H | X \rangle$ is still the energy of a classical lattice gas in which the equivalence of the canonical and microcanonical ensembles implies the analogue of Eq. (10) and, via the argument of section 4, the nonuniformity of the ground state.

The real novelty introduced by the additional interactions is the possible coexistence of classical long-range order with the off-diagonal one. A thorough discussion of this question, concluding negatively, can be found already in Penrose and Onsager's 1956 paper. Nevertheless, a controversy has remained until recently, when reliable quantum Monte Carlo computations on square lattice models have shown such a coexistence (Batrouni *et al* 1995, Scalettar *et al* 1995). We hope that the ideas developed in this paper can contribute in the future to further elucidate this interesting problem.

This work was supported by the Hungarian Scientific Research Fund (OTKA) under Grant No. T 30543.

Appendix. A digression on ODLRO

Following the standard definition (Penrose and Onsager 1956, PO), we give below the expression of the ODLRO parameter in terms of (a_X) , valid for any Bose gas whose Hamiltonian contains a hard-core repulsion and conserves the number of particles. The density operator in the ground state Ψ_0 (supposed to be normalized) is the orthogonal projection $|\Psi_0\rangle\langle\Psi_0|$. The density matrix in the basis (4) is therefore $(a_X a_Y)$. From here the one-particle reduced density matrix $\sigma = (\sigma_{xy})$ is obtained by taking a partial trace over $N - 1$ particle positions,

$$\sigma_{xy} = \sum_{X'} a_{X' \cup \{x\}} a_{X' \cup \{y\}} . \quad (96)$$

In the above sum X' runs over the $N - 1$ -point subsets of Λ not containing x and y . The matrix σ is real symmetric, positive semidefinite (in fact, positive definite for $N > 2$), its trace is N and all its elements are positive. A possible choice for the order parameter, according to PO, is

$$\omega = |\Lambda|^{-2} \sum_{x, y \in \Lambda} \sigma_{xy} . \quad (97)$$

Another definition of PO for the order parameter is $1/|\Lambda|$ times the largest eigenvalue of σ . If Λ is a shift-invariant set (which supposes periodic boundary conditions), the ground state is translation invariant and

$\sigma_{xy} = \sigma_{x-y}$. Applying the Perron-Frobenius theorem to σ we find that its largest eigenvalue is $\sum_y \sigma_{x-y} = |\Lambda|\omega$, so that the above two definitions coincide. The order parameter ω also has an interpretation as the density of the Bose-Einstein condensate. This holds because the operator associated with the density of the condensate is

$$|\Lambda|^{-1} b_{k=0}^* b_{k=0} = |\Lambda|^{-2} \sum_{x,y \in \Lambda} b_x^* b_y \quad (98)$$

and because

$$\langle \Psi_0 | b_x^* b_y | \Psi_0 \rangle = \sigma_{xy} . \quad (99)$$

Inserting (96) into (97) we obtain

$$\omega = \omega' + \frac{N}{|\Lambda|^2} \quad \omega' = \frac{1}{|\Lambda|^2} \sum_X a_X \sum_{x \in X} \sum_{y \notin X} a_{X \cup \{y\} \setminus \{x\}} . \quad (100)$$

The maximum of ω' under the condition of normalization is $\rho(1 - \rho)$, attained with a_X constant. It is realized by H_0 on complete graphs and by the ferromagnetic Heisenberg model on arbitrary graphs. Our ω' is one fourth of the order parameter usually used in the spin- $\frac{1}{2}$ XY model, cf. Fujiki and Betts 1986, Hamer *et al* 1999. There is ODLRO for a density ρ if ω or, equivalently, ω' has a nonvanishing limit $\omega_0 > 0$ as N and $|\Lambda|$ go to infinity while $N/|\Lambda|$ goes to ρ . When this occurs, conservation of the particle number is spontaneously broken and there appears (at least) a one-parameter continuum of ground states in infinite volume. The order parameter operator, $\sum b_x^*$, is nondiagonal in the natural basis (4) (whence the name ODLRO). The ground states γ_α , $0 \leq \alpha < 2\pi$, are symmetry-related. For any local operator B

$$\gamma_\alpha(B) = \gamma_0 \left(\prod_x e^{-i\alpha n_x} B \prod_x e^{i\alpha n_x} \right) . \quad (101)$$

In particular,

$$\gamma_\alpha(b_x^*) = e^{-i\alpha} \gamma_0(b_x^*) \neq 0 . \quad (102)$$

References

- [1] Batrouni G G, Scalettar R T, Zimanyi G T and Kampf A P 1995
Phys. Rev. Letters **74** 2527
Scalettar R T, Batrouni G G, Kampf A P and Zimanyi G T 1995
Phys. Rev. B **51** 8467
- [2] Biggs N 1974 *Algebraic Graph Theory* (Cambridge University Press)
- [3] Dobrushin R L and Shlosman S B 1994 *Adv. Sov. Math.* **20** 91
- [4] Dyson F J, Lieb E H and Simon B 1978 *J. Stat. Phys.* **18** 335
- [5] Fujiki S and Betts D D 1986 *Can J. Phys.* **64** 876
- [6] Hamer C J, Hövelborn T and Bachhuber M 1999
J. Phys. A: Math. Gen. **32** 51
- [7] Kennedy T, Lieb E H and Shastry B S 1988 *Phys. Rev. Letters* **61** 2583
- [8] Kubo K and Kishi T 1988 *Phys. Rev. Lett.* **61** 2585
- [9] Lewis J T, Pfister C E and Sullivan W G 1994 *J.Stat. Phys.* **77** 397
- [10] Matsubara T and Matsuda H 1956 *Prog. Theor. Phys.* **16** 569
- [11] Penrose O 1991 *J. Stat. Phys.* **63** 761
- [12] Penrose O and Onsager L 1956 *Phys. Rev.* **104** 576
- [13] Pfister C E 1991 *Helv. Phys. Acta* **64** 953
- [14] Pleimling M and Hüller A 2000 *Preprint* cond-mat/0011116
- [15] Tóth B 1991 *J. Stat. Phys.* **65** 373
- [16] Tóth B 1990 *J. Stat. Phys.* **61** 749
- [17] Yang C N 1962 *Rev. Mod. Phys.* **34** 694
- [18] Zhang S and Runge K 1992 *Phys. Rev. B* **45** 1052